## AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior listings of claims in the application.

1. (currently amended) A compound of the formula (I):



(I)

wherein,

each $R^1$, $R^2$, and $R^3$ are independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

X is N~~, O, or S~~;

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, $CH_2OH$, $CH(OH)CH{=}CH_2$, or $C(O)NHCHR^{10}CO_2H$;

each $R^5$ is independently H, alkyl, alkenyl, aryl, heteroaryl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^2$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{10}$ is independently an amino acid side chain;

each $P^1$ and $P^2$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group;

or pharmaceutically acceptable salts thereof.

2. (currently amended) The compound of claim 1, wherein:

X is N ~~or O~~;

$R^1$ is alkyl substituted with aryl, which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

$R^4$ is H, $CON(R^7)_2$, $C(O)NHCHR^{10}CO_2H$, or $CH_2OH$;

each $R^5$ is independently H, alkyl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^2$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{10}$ is independently an amino acid side chain;

each $P^1$ and $P^2$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group.

3. (currently amended) The compound of claim 1, wherein:

X is N ~~or O~~;

$R^1$ is alkyl substituted with aryl, which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, halo, or $C_{1-6}$ alkyl;

$R^4$ is H, $CONHR^7$, or $CH_2OH$;

each $R^5$ is independently H or alkyl;

each $R^6$ is independently H or alkyl;

$R^7$ is H, alkyl, or $P^2$; and

$P^2$ is a nitrogen protecting group.

4. (currently amended) The compound of claim 1, wherein:

X is N ~~or O~~;

$R^1$ is alkyl substituted with aryl, which is optionally substituted with 1-5 substituents selected from OH or $C_{1-6}$ alkyl; and

$R_4$ is H, $CONH_2$, or $CH_2OH$.

5. (currently amended) The compound of claim 1, wherein:

X is N ~~or O~~;

$R^1$ is $C_1$ alkyl substituted with phenyl, which is substituted at the 2- and 6- positions with Me and is substituted at the 4- position with OH; and

$R^4$ is H, $CONH_2$, or $CH_2OH$.

6. (currently amended) The compound of claim 1 having the formula (II):




(II)

wherein,

X is N ~~or O~~;

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, $CH_2OH$, or $C(O)NHCHR^{10}CO_2H$;

each $R^5$ is independently H, alkyl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^2$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^9$ is independently $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

each $R^{10}$ is independently an amino acid side chain;

each n is independently 0, 1, 2, 3, 4, or 5;

each $P^1$ and $P^2$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group.

7. (original) The compound of claim 6, wherein:

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, or $CH_2OH$;

each $R^5$ is independently H, alkyl, or acyl;

each $R^6$ is independently H or alkyl;

each $R^7$ is independently H or alkyl;

each $R^9$ is independently $OR^6$, CN, $NO_2$, halo, or $C_{1-6}$ alkyl; and

each n is independently 0 1, 2, or 3.

8. (original) The compound of claim 6, wherein:

$P^1$ is a BOC or Fmoc;

$P^2$ is a solid support; and

$P^3$ is *t*-Bu, Bn, Me, or Ac.

9. (original) The compound of claim 6, wherein:

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, or $CH_2OH$;

each $R^5$ is independently H, alkyl, acyl, or $P^1$;

each $R^6$ is independently H or $P^3$;

each $R^7$ is independently H or $P^2$;

each $R^9$ is independently $OR^6$ or $C_{1-6}$ alkyl;

each n is independently 0, 1, or 2;

$P^1$ is a BOC;

$P^2$ is a solid support; and

$P^3$ is *t*-Bu.

10. (original) The compound of claim 6, wherein:

$R^4$ is H, $CONH_2$, or $CH_2OH$;

each $R^5$ is independently H, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is H or alkyl

each $R^9$ is $C_{1-6}$ alkyl or $OR^6$;

each $R^{10}$ is independently an amino acid side chain;

each n is independently 1, 2, or 3; and

$P^1$ is a nitrogen protecting group.

11. (currently amended) The compound of claim 1 that is formula (III):



(III)

wherein,

X is ~~O or~~ N;

$R^9$ is $C_{1-6}$ alkyl; and

n is 2.

12. (currently amended) The compound of claim 1 that is formula (IV):



(IV)

wherein X is N ~~or O~~.

13. (currently amended) The compound of claim 1 having the formula (V):




(S, S, S, R)

(V)

wherein

X is N ~~or O~~; and

$R^4$ is $CONH_2$, H, or $CH_2OH$.

14. (original) The compound of claim 1 having the formula (VI):




(VI)

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, $CH_2OH$, or $CH(OH)CH{=}CH_2$, or $C(O)NHCHR^{10}CO_2H$;

each $R^5$ is independently H, alkyl, alkene, aryl, heteroaryl, acyl, or $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^2$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{10}$ is independently an amino acid side chain;

each $P^1$ and $P^2$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group.

15. (original) The compound of claim 14, wherein:

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, halo, or $C_{1-6}$ alkyl;

$R^4$ is H, $CON(R^7)_2$, or $CONHR^7$, or $C(O)NHCHR^{10}CO_2H$;

each $R^5$ is independently H, alkyl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$; and

each $R^{10}$ is independently an amino acid side chain.

16. (original) A method of making a compound of the formula (VIII):



(VIII)

comprising coupling compounds of the formulas (XI) and (XII)



(XI)



(XII)

using a ruthenium catalyst, to give a compound of formula (IX); and



(IX)

reacting the compound of formula (IX) with a deprotecting agent to give a compound of the formula (VIII);

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^4$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{11}$, $R^{12}$, and $R^{13}$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^{16}$, CN, $NO_2$, $NHR^{17}$, $N(R^{17})_2$, halo, $CONHR^{17}$, $CON(R^{17})_2$, $CO_2R^{18}$, or $C_{1-6}$ alkyl;

each $R^{16}$ is independently H, alkyl, or $P^3$;

each $R^{17}$ is independently H, alkyl, acyl, or $P^4$;

each $R^{18}$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $P^1$, $P^2$, and $P^4$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group.

17. (original) A method of making a compound of the formula (XVI):



(XVI)

comprising coupling compounds of the formulas (XI) and (XIII)



(XI)



(XIII)

by first reacting the free alcohols with a silicon protecting group, and then treating the resulting compound with a ruthenium catalyst, giving a compound of the formula (VII);



(VII)

reacting the compound of formula (VII) under pH conditions sufficient to remove acid labile protecting groups, if any;

treating the resulting product with peroxide and base under conditions sufficient to hydrolyze the thioester; and

coupling the resulting product with a solid phase peptide, giving a compound of the formula (XVI);

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^4$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{11}$ and $R^{12}$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^{16}$, CN, $NO_2$, $NHR^{17}$, $N(R^{17})_2$, halo, $CONHR^{17}$, $CON(R^{17})_2$ $CO_2R^{18}$, or $C_{1-6}$ alkyl;

each $R^{16}$ is independently H, alkyl, or $P^3$;

each $R^{17}$ is independently H, alkyl, acyl, or $P^4$;

each $R^{18}$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $P^1$ and $P^4$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group; and

$P^5$ is a sulfur protecting group.

18. (currently amended) A method of making a compound of the formula (XIV):



(XIV)

comprising coupling compounds of formulas (XI) and (XIII),



(XI)



(XIII)

with a ruthenium catalyst;

treating the resulting compound with peroxide and base under conditions sufficient to hydrolyze the thioester;

amidation or esterification of the resulting acid; and

treatment of the resulting compound with a deprotecting agent sufficient to remove protecting groups, giving a compound of the formula (XIV);

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

X is N ~~or O~~;

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, $CH_2OH$, or $CH(OH)CH=CH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^4$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{11}$ and $R^{12}$ are independently alkyl substituted with aryl or heteroaryl, each of

which is optionally substituted with 1-5 substituents selected from $OR^{16}$, CN, $NO_2$, $NHR^{17}$, $N(R^{17})_2$, halo, $CONHR^{17}$, $CON(R^{17})_2$, $CO_2R^{18}$, or $C_{1-6}$ alkyl;

each $R^{16}$ is independently H, alkyl, or $P^3$;

each $R^{17}$ is independently H, alkyl, acyl, or $P^4$;

each $R^{18}$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $P^1$ and $P^4$ is independently a nitrogen protecting group;

each $P^3$ is independently an oxygen protecting group; and

$P^5$ is a sulfur protecting group.

19. (original) A method of making a compound of formula (XVII):



(XVII)

comprising coupling compounds of formulas (XI) and (XIII)



(XI)



(XIII)

with a ruthenium catalyst;

treating the resulting compound with peroxide and base under conditions sufficient to hydrolyze the thioester; and

reacting the free hydroxyls with an oxygen protecting group to give a compound of formula (XVIII)



(XVIII)

coupling the compound of formula (XVIII) with an alcohol of formula $R^{13}(CHOH)CHOR^{16}$; and

treating the resulting compound with a deprotecting agent sufficient to remove protecting groups to give a compound of formula (XVII);

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^4$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{11}$, $R^{12}$, and $R^{13}$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^{16}$, CN, $NO_2$, $NHR^{17}$, $N(R^{17})_2$, halo, $CONHR^{17}$, $CON(R^{17})_2$ $CO_2R^{18}$, or $C_{1-6}$ alkyl;

each $R^{16}$ is independently H, alkyl, or $P^3$;

each $R^{17}$ is independently H, alkyl, acyl, or $P^4$;

each $R^{18}$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $P^1$ and $P^4$ is independently a nitrogen protecting group;

each $P^3$ is independently an oxygen protecting group; and

$P^5$ is a sulfur protecting group.

20. (original) A composition comprising a compound of formula (I) in claim 1 and a pharmaceutically acceptable carrier.

21. (currently amended) A compound of formula (XIX):



(XIX)

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$ halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

X is N~~, O, or S~~;

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, $CH_2OH$, $CH(OH)CH{=}CH_2$, or $C(O)NHCHR^{10}CO_2H$;

each $R^5$ is independently H, alkyl, alkenyl, aryl, heteroaryl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^2$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{10}$ is independently an amino acid side chain;

each $P^1$ and $P^2$ is independently a nitrogen protecting group;

each $P^3$ is independently an oxygen protecting group; and

or pharmaceutically acceptable salts thereof.

22. (currently amended) The compound of claim 21 wherein:

X is N or ~~O~~;

$R^1$ is alkyl substituted with aryl, which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

$R^4$ is H, $CON(R^7)_2$, $C(O)NHCHR^{10}CO_2H$, or $CH_2OH$; and

each $R^5$ is independently H, alkyl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^{10}$ is independently an amino acid side chain.

23. (currently amended) The compound of claim 21, wherein:

X is N ~~or O~~;

$R^1$ is alkyl substituted with aryl, which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, halo, or $C_{1-6}$ alkyl;

$R^4$ is H, $CONHR^7$, or $CH_2OH$;

each $R^5$ is independently H or alkyl;

each $R^6$ is independently H or alkyl; and

$R^7$ is H, alkyl, or $P^2$.

24. (currently amended) The compound of claim 21, wherein:

X is N ~~or O~~;

$R^1$ is alkyl substituted with aryl, which is optionally substituted with 1-5 substituents selected from OH or $C_{1-6}$ alkyl; and

$R^4$ is H, $CONH_2$, or $CH_2OH$.

25. (currently amended) The compound of claim 21, wherein:

X is N ~~or O~~;

$R^1$ is $C_1$ alkyl substituted with phenyl, which is substituted at the 2- and 6- positions with Me and is substituted at the 4- position with OH; and

$R^4$ is H, $CONH_2$, or $CH_2OH$.

26. (original) The compound of claim 21, wherein

X is N;

$R^1$ is methyl substituted with phenyl, which is substituted at the 4- position with OH; and

$R^4$ is $CONH_2$.

27. (currently amended) The compound of claim 21 having the formula (XX):




(XX)

wherein,

X is N ~~or O~~;

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, $CH_2OH$, or $C(O)NHCHR^{10}CO_2H$;

each $R^5$ is independently H, alkyl, acyl, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl; or $P^2$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^9$ is independently $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

each $R^{10}$ is independently an amino acid side chain;

each n is independently 0, 1, 2, 3, 4, or 5;

each $P^1$ and $P^2$ is independently a nitrogen protecting group; and

each $P^3$ is independently an oxygen protecting group.

28. (original) The compound of claim 27, wherein:

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, or $CH_2OH$;

each $R^5$ is independently H, alkyl, or acyl;

each $R^6$ is independently H or alkyl;

each $R^7$ is independently H or alkyl;

each $R^9$ is independently $OR^6$, CN, $NO_2$, halo, or $C_{1-6}$ alkyl; and

each n is independently 0 1, 2, or 3.

29. (original) The compound of claim 27, wherein:

$R^4$ is H, $CON(R^7)_2$, $CONHR^7$, or $CH_2OH$;

each $R^5$ is independently H, alkyl, acyl, or $P^1$;

each $R^6$ is independently H or $P^3$;

each $R^7$ is independently H or $P^2$;

each $R^9$ is independently $OR^6$ or $C_{1-6}$ alkyl;

each n is independently 0 or 1;

$P^1$ is a BOC;

$P^2$ is a solid support; and

$P^3$ is *t*-Bu.

30. (original) The compound of claim 27, wherein:

$R^4$ is H, $CONH_2$, or $CH_2OH$;

each $R^5$ is independently H, $P^1$, or $C(O)CHR^{10}NH_2$;

each $R^6$ is H or alkyl

each $R^9$ is $C_{1-6}$ alkyl or $OR^6$;

each $R^{10}$ is independently an amino acid side chain;

each n is independently 1, 2, or 3; and

$P^1$ is a nitrogen protecting group.

31. (currently amended) The compound of claim 21 having the formula (XXI):




(XXI)

wherein,

X is ~~O or~~ N;

$R^4$ is H, $CONH_2$, or $CH_2OH$;

$R^9$ is $C_{1-6}$ alkyl; and

n is 2.

32. (original) The compound of claim 21 having the formula (XXII):



(S, S, R, S)

(XXII)

wherein

$R^2$ is $C_{1-6}$ alkyl; and

n is 0, 1, or 2.

33. (original) A method of making a compound of formula (XXIII):



(XXIII)

comprising coupling compounds of formulas (XXV) and (XIII)



(XXV)



(XIII)

using a ruthenium catalyst, giving a compound of the formula (XXIV);

treating the resulting product with peroxide and base under conditions sufficient to hydrolyze the thioester;

coupling the resulting product with a solid phase peptide; and

treating the resulting compound with a deprotecting agent, giving a compound of the formula (XXIII);

wherein,

each $R^1$, $R^2$, and $R^3$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^6$, CN, $NO_2$, $NHR^7$, $N(R^7)_2$, halo, $CONHR^7$, $CON(R^7)_2$, $CO_2R^8$, or $C_{1-6}$ alkyl;

each $R^6$ is independently H, alkyl, or $P^3$;

each $R^7$ is independently H, alkyl, acyl, or $P^4$;

each $R^8$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $R^{11}$ and $R^{12}$ is independently alkyl substituted with aryl or heteroaryl, each of which is optionally substituted with 1-5 substituents selected from $OR^{16}$, CN, $NO_2$, $NHR^{17}$, $N(R^{17})_2$, halo, $CONHR^{17}$, $CON(R^{17})_2$, $CO_2R^{18}$, or $C_{1-6}$ alkyl;

each $R^{16}$ is independently H, alkyl, or $P^3$;

each $R^{17}$ is independently H, alkyl, acyl, or $P^4$;

each $R^{18}$ is independently H, alkyl, aralkyl, or heteroaralkyl;

each $P^1$ and $P^4$ is independently a nitrogen protecting group;

each $P^3$ is independently an oxygen protecting group; and

$P^5$ is a sulfur protecting group.

34. (original) A composition comprising a compound of formula (XIX) in claim 21 or pharmaceutically acceptable salts thereof and a pharmaceutically acceptable carrier.

35. (currently amended) A method of treating a mu opioid receptor (MOR) mediated disorder that is pain in a subject comprising administering a compound of formula (I) in claim 1 or a compound of formula (XIX) in claim 21 or pharmaceutically acceptable salts thereof.

36. (currently amended) A method of treating a mu opioid receptor (MOR) mediated disorder that is pain in a subject comprising administering a composition comprising a compound of formula (I) in claim 1 or a compound of formula (XIX) in claim 21 or pharmaceutically acceptable salts thereof.

37. (original) A method of treating pain in a subject, comprising administering to the subject a compound of formula (I) in claim 1 or of formula (XIX) in claim 21 or pharmaceutically acceptable salts thereof.

38. (cancelled)